

Negative Probability and Uncertainty Relations

Thomas Curtright*

Department of Physics, University of Miami, Box 248046, Coral Gables, Florida 33124, USA

Cosmas Zachos†

High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4815, USA

(Dated: May 23, 2001)

A concise derivation of all uncertainty relations is given entirely within the context of phase-space quantization, without recourse to operator methods, to the direct use of Weyl's correspondence, or to marginal distributions of x and p .

PACS numbers: 03.65.Ta, 03.65.Db, 03.65.Fd, 11.15.Kc

Phase-space quantization is the third autonomous and logically complete formulation of quantum mechanics beyond the conventional ones based on operators in Hilbert space or path integrals [1, 2, 3]. It is free of operators and wavefunctions: observables and matrix elements are computed through phase-space integrals of c-number functions ("classical kernels") weighted by a Wigner function (WF) [3, 4]. This is a phase-space distribution function which is *not* positive semi-definite, and constitutes the Weyl correspondent [5] of the density matrix in the conventional formulation,

$$f_{mn}(x, p) = \frac{1}{2\pi} \int dy e^{-iyp} \psi_m^*(x - \frac{\hbar}{2}y) \psi_n(x + \frac{\hbar}{2}y) = f_{nm}^*(x, p). \quad (1)$$

Operators of the conventional formulation, when properly ordered (eg, Weyl-ordered), correspond uniquely to phase-space classical kernel functions, while operator products correspond to $\star$ -products [6] of these classical kernels, the $\star$ -product being a noncommutative and associative operation encoding quantum mechanical action. The above wavefunctions, however, may be forfeited, since the WFs are determined, in principle, as the solutions of the celebrated $\star$ -genvalue functional equations [7, 8, 9, 10]. Connections to the original, operator, formulation of quantum mechanics may thus be ignored.

Recent M-theory advances linked to noncommutative geometry and matrix models [11] apply spacetime uncertainty principles [12] reliant on phase-space quantization and the $\star$ -product. Transverse spatial dimensions act formally as momenta, and, analogously to quantum mechanics, their uncertainty is increased or decreased inversely to the uncertainty of a given direction.

For classical (non-negative) probability distributions, expectation values of non-negative functions are likewise non-negative, and thus result in standard constraint inequalities for the constituent pieces of such functions. On the other hand, in phase-space quantization, the distribution functions are *non-positive-definite*, such as, in general, the *quasi*-probability WF: it was interpreted early on by Bartlett [13], and later by Feynman [14], as a "negative probability function", with the proper non-negative marginal probabilities upon projection to either x or p space. Hence, a frequent first question in phase-space quantization is how Heisenberg's standard quantum mechanical uncertainty relation arises for moments of such distributions.

To be sure, Moyal derived these uncertainty relations, in his original formulation of quantum mechanics in phase space, by careful analysis of conditioned and marginal probabilities. Nevertheless, plain evaluations of expectation values of the c-number variables $\langle x^2 \rangle$, $\langle p^2 \rangle$, etc, do not evince constraints; and the student of deformation quantization is left wondering how $\hbar$ enters the constraint of such expectation values of (c-number) observables when the variables x, p do not contain $\hbar$. How do their moments manage to constrain each other by extracting $\hbar$ out of the Wigner function?

The answer lies in Groenewold's associative $\star$ -product [6],

$$\star \equiv e^{i\hbar(\overleftarrow{\partial}_x \overrightarrow{\partial}_p - \overleftarrow{\partial}_p \overrightarrow{\partial}_x)/2}, \quad (2)$$

which is the cornerstone of phase-space quantization. Its mechanics is reviewed in [7, 10, 15]. An alternate, integral,

*Electronic address: curtright@physics.miami.edu

†Electronic address: zachos@hep.anl.gov

representation of this product is [16]

$$f \star g = (\hbar\pi)^{-2} \int dudvdwdz f(x+u, p+v) g(x+w, p+z) \exp\left(\frac{2i}{\hbar}(uz - vw)\right), \quad (3)$$

which readily displays associativity. The phase-space trace is directly seen in this representation to obey

$$\int dpdx f \star g = \int dpdx fg = \int dpdx g \star f. \quad (4)$$

The WF spectral properties [1] are reviewed and illustrated in [8, 10]. Eg, the trace-normalization condition,

$$\int dx dp f_{mn}(x, p) = \delta_{mn}, \quad (5)$$

and the spectral orthogonality conditions [7], $f_{mn} \star f_{kl} = \delta_{ml} f_{kn}/2\pi\hbar$. Given (4), it follows that $\int dx dp f_{mn}(x, p) f_{lk}^*(x, p) = \delta_{ml} \delta_{nk}/2\pi\hbar$. For complete sets of input wavefunctions, it also follows that

$$\sum_{m,n} f_{mn}(x, p) f_{mn}^*(x', p') = \frac{1}{2\pi\hbar} \delta(x - x') \delta(p - p'). \quad (6)$$

An arbitrary phase-space function $\varphi(x, p)$ can thus be expanded as $\varphi(x, p) = \sum_{m,n} c_{mn} f_{mn}(x, p)$.

Here, a concise proof of all uncertainty relations is provided completely within the autonomous framework of phase-space quantization, unlike extant discussions of such correlation inequalities, which rely on the operator formulation of quantum mechanics. It is stressed that, in the following, *no* operators occur, only the $\star$ -product operation, and x and p are c-numbers. The controlling fact is that expectation values of arbitrary real $\star$ -squares are positive semi-definite, even though the Wigner distribution $f(x, p)$ itself is not. Specifically, for any complex phase-space function $g(x, p)$, and any (real) Wigner function $f(x, p)$ representing a pure state, the following inequality holds:

$$\langle g^* \star g \rangle = \int dpdx (g^* \star g) f \geq 0. \quad (7)$$

The $\star$ is absolutely crucial here, and its removal leads to violation of the inequality, as can easily be arranged by choosing the support of g to lie mostly in those regions of phase-space where the Wigner function is negative. (The only pure state WF which is non-negative is the Gaussian [3, 8, 17]). In Hilbert space operator formalism, this relation (7) would correspond to the positivity of the norm. By (4), $\int dpdx (g^* \star g) f = \int dpdx (g^* \star g) \star f$, ie inside a phase-space integral an ordinary product can be extended to a $\star$ -product, provided it not be part of a longer string. Eg, the one $\star$ -product of the left hand side cannot be eliminated, because of the extra ordinary product with f .

To prove the inequality (7), it suffices to recognize that, for a pure state, its (real) Wigner function can be expanded in a complete basis of Wigner $\star$ -genfunctions of a convenient Hamiltonian, [10], $f = \sum_{n,m} c_m^* c_n f_{mn}$, for complex coefficients c_n , s.t. $\sum_n |c_n|^2 = 1$, to satisfy (5). Then, it follows that [18, 19]

$$f \star f = f/h. \quad (8)$$

Consequently, given the relations (4), $(g \star f)^* = f \star g^*$, and the associativity of the $\star$ -product,

$$\int dpdx (g^* \star g) f = h \int dx dp (g^* \star g) (f \star f) = h \int dx dp (f \star g^*) \star (g \star f) = h \int dx dp |g \star f|^2. \quad (9)$$

This expression, then, involves a real non-negative integrand and is itself positive semi-definite. (Similarly, if f_1 and f_2 are pure state WFs, the transition probability between the respective states [3] is also manifestly non-negative by the same argument: $\int dpdx f_1 f_2 = (2\pi\hbar)^2 \int dx dp |f_1 \star f_2|^2$.)

Given (7), correlations of observables follow conventionally from specific choices of $g(x, p)$. For example, to produce Heisenberg's uncertainty relation, one only need choose

$$g = a + bx + cp, \quad (10)$$

for arbitrary complex coefficients a, b, c . The resulting positive semi-definite quadratic form is then

$$a^* a + b^* b \langle x \star x \rangle + c^* c \langle p \star p \rangle + (a^* b + b^* a) \langle x \rangle + (a^* c + c^* a) \langle p \rangle + c^* b \langle p \star x \rangle + b^* c \langle x \star p \rangle \geq 0, \quad (11)$$

for any a, b, c . The eigenvalues of the corresponding matrix are then non-negative, and thus so must be its determinant. Given

$$x \star x = x^2, \quad p \star p = p^2, \quad p \star x = px - i\hbar/2, \quad x \star p = px + i\hbar/2, \quad (12)$$

and the usual

$$(\Delta x)^2 \equiv \langle (x - \langle x \rangle)^2 \rangle, \quad (\Delta p)^2 \equiv \langle (p - \langle p \rangle)^2 \rangle, \quad (13)$$

this condition on the 3×3 matrix determinant amounts to

$$(\Delta x)^2 (\Delta p)^2 \geq \hbar^2/4 + \left(\langle (x - \langle x \rangle)(p - \langle p \rangle) \rangle \right)^2, \quad (14)$$

and hence

$$\Delta x \Delta p \geq \hbar/2. \quad (15)$$

The inequality is saturated for a vanishing original integrand $g \star f = 0$, for suitable a, b, c , and when the last term of (14) vanishes: x, p statistical independence, such as in a Gaussian ground state WF, $f_{00} = 2h \exp(-(x^2 + p^2)/h)$.

More general choices of g will likewise constrain as many observables as this function has terms (-1 , if there is a constant term). For instance, for more general (real) observables $u(x, p), v(x, p)$, the resulting inequality is

$$\Delta u \Delta v \geq \frac{1}{2} \sqrt{|\langle u \star v - v \star u \rangle|^2 + \langle (u - \langle u \rangle) \star (v - \langle v \rangle) + (v - \langle v \rangle) \star (u - \langle u \rangle) \rangle^2}. \quad (16)$$

The minimum uncertainty is realized at $\langle u \star v + v \star u \rangle = 2\langle u \rangle \langle v \rangle$, with $g \star f = 0$ for specific coefficients, ie,

$$\left(\Delta u (v - \langle v \rangle) - ki \Delta v (u - \langle u \rangle) \right) \star f = 0, \quad (17)$$

where k is the sign of $i\langle u \star v - v \star u \rangle$. Solving such $\star$ -equations is elaborated in [8, 9, 10, 15].

This work was supported in part by the US Department of Energy, Division of High Energy Physics, Contract W-31-109-ENG-38, and the NSF Award 0073390.

-
- [1] J Moyal, Proc Camb Phil Soc **45** (1949) 99-124
 - [2] F Bayen, M Flato, C Fronsdal, A Lichnerowicz, and D Sternheimer, Ann Phys **111** (1978) 61; *ibid.* 111;
 - [3] For reviews, see M Hillery, R O'Connell, M Scully, and E Wigner, Phys Repts **106** (1984) 121;
H-W Lee, Phys Repts **259** (1995) 147;
N Balasz and B Jennings, Phys Repts **104** (1984) 347;
R Littlejohn, Phys Repts **138** (1986) 193;
M Berry, Philos Trans R Soc London A287 (1977) 237;
M Gadella, Fortschr Phys **43** (1995) 3, 229-264;
L Cohen, *Time-Frequency Analysis* (Prentice Hall PTR, Englewood Cliffs, 1995);
F Berezin, Sov Phys Usp **23** (1980) 763-787;
P Carruthers and F Zachariasen, Rev Mod Phys **55** (1983) 245;
A M Ozorio de Almeida, Phys Rept **295** (1998) 265-342
 - [4] E Wigner, Phys Rev **40** (1932) 749-759
 - [5] H Weyl, Z Phys **46** (1927) 1; also reviewed in H Weyl (1931) *The Theory of Groups and Quantum Mechanics*, Dover, New York
 - [6] H Groenewold, Physica **12** (1946) 405-460
 - [7] D Fairlie, Proc Camb Phil Soc **60** (1964) 581-586
 - [8] T Curtright, D Fairlie, and C Zachos, Phys Rev **D58** (1998) 025002
 - [9] M Hug, C Menke, and W Schleich, Phys Rev **A57** (1998) 3188-3205
 - [10] T Curtright, T Uematsu, and C Zachos, J Math Phys **42** (2001) 2396-2415 [hep-th/0011137]
 - [11] N Seiberg and E Witten, JHEP **9909** (1999) 032;
for a review, see L Castellani, Class Quant Grav **17** (2000) 3377-3402 [hep-th/0005210]
 - [12] T Yoneya, Mod Phys Lett **A4** (1989) 1587;
A Jevicki and T Yoneya, Nucl Phys **B535** (1998) 335 [hep-th/9805069];
N Seiberg, L Susskind and N Toumbas, JHEP **0006** (2000) 044 [hep-th/0005015]

- [13] M Bartlett, Proc Camb Phil Soc **41** (1945) 71-73
- [14] R Feynman, "Negative Probability" in *Essays in Honor of David Bohm*, B Hiley and F Peat (eds) (Routledge and Kegan Paul, London, 1987); for a popular review, see D Leibfried, T Pfau, and C Monroe, Physics Today, April 1998, 22-28
- [15] F Hansen, Rept Math Phys **19** (1984) 361-381;
C Roger and V Ovsienko, Russ Math Surv **47** (1992) 135-191;
C Zachos, J Math Phys **41** (2000) 5129-5134 [hep-th/9912238]
- [16] J v Neumann, Math Ann **104** (1931) 570-578
- [17] R Hudson, Rep Math Phys **6** (1974) 249-252
- [18] T Takabayasi, Prog Theo Phys **11** (1954) 341-373
- [19] G Baker, Phys Rev **109** (1958) 2198-2206